INDEX TO SCIENTIFIC NAMES

HERPETOLOGICA

Volume 43 December 1987

Acacia 241
Acer saccharum 329
Aeluroglena 202
Agama agama 74
Agkistrodon 101, 239
Ahaetulla 246
Alligator mississippiensis 255–256
Alocasia odora 483
Alopoglossus angulatus 386
  atriventris 386
  carinicaudatus 386
Alsophis 241
  cantherigerus 247
  spp. 246
  vudii 246
Amblyrhynchus cristatus 312
Ambystoma maculatum 496–500
  opacum 496–500
  tigrinum 500
  tremblayi 106, 111, 115
Ameiva ameiva 386
  chrysolaema 241, 243–247
  exsul 374–375, 377–382
  quadrilineata 380
  sp. 243
  taeniura 243
Amolops afghanus 95, 97
  chunganensis 97
  formosus 97
  hainanensis 97
  kinabaluensis 97
  ***macrorhynchus* 96**–97
  nasica 97
  ricketti 97
Amphibolurus muricatus 236
Amphiuma means 113, 492
  pholeter 106–107, 113, 115–116
  tridactylum 106, 113, 115
Anadia ***altaserrania*** 183–184, **186**–189
  bitaeniata 182–190
  bogotensis 188–189
  brevifrontalis 182–184, 186, 188–190
  pamplonensis 182–185, 188–190
  pulchella 182–185, 188–190
  pulchra 184
Anas junius 371
Anax junius 370–371
Anolis 241
  bonairensis 312
  carolinensis 34
  caudalis 243
  chlorocyanus 243, 245, 247
  coelestinus 243–245, 247
  cristatellus 132
  cybotes 243–245, 247
  distichus 243–245, 247
  longiceps 132
  olssoni 243–244, 246–247
  punctatus 386
  punctatus boulengeri 386
  semilineatus 243–245
  sp. 243
  wattsi 132
Antechinus swainsoni 236
Ardisia sieboldii 483
Argyrogena 203
Aristida stricta 421
Arizona 201–202, 204–206
  elegans 203
Ascaphus 128
Atheris squamigera 101
Atractus flammigerus 387
  flammigerus flammigerus 387
Austrelaps sp. 35
Bachia vermiforme 386
  vermiformis 386
Barisia 35
  imbricata 29–30, 33–34, 36
  imbricata ciliaris 30, 32–34
  imbricata imbricata 29–31, 33–34, 36
  imbricata jonsei 30, 33–34
  imbricata planifrons 30, 33–34
  monticola 29, 34
Batrachuperus pinchoni 115
Bellamya javanica 5
Belostoma sp. 370–371
Betula pendula 406
Bishofia javanica 483
Boa hortulana 101
Bogabdella diversa 303
Bombina orientalis 384
Bothrops asper 133
  atrox 133
  bilineatus 386
  bilineatus bilineatus 386
  bilineatus smaragdina 386
  castelnaudi 387
  hyoprora 388
  nigroviridis aurifer 294
  taeniatus 387
Bufo 212
  americanus 214, 369–371, 478
  boreas 482
  bufo 221, 478
  calamita 478
  melanostictus 1
Caiman crocodilus 256
  yacare 255
Calluna vulgaris 346–347
Canavalia obtusifolia 319
Caretta caretta 79, 307–313, 315
Caudiverbera 213
Celestus 131
Cemophora 201–202

Cerdocyon thous 255
Chalcides chalcides 35
ocellatus 312
Chamaelinorops barbouri 132
Chelonia 130–131
mydas 74–76, 308, 310–312, 315
Chelus fimbriatus 387
Chelydra serpentina 39–40, 42–50, 52, 324, 393
serpentina osceola 326
Chironeus cochranae 387
multiventris 387
Chlorophilus olivaceus 163
Chrysemys 88
picta 40, 50, 52, 55, 82, 87, 305, 312–313, 326, 334, 393
picta belli 82
picta marginata 55–56, 61–62
scripta 239
Cinnamomum japonicum 483
Circus approximans 342
Clelia bicolor 387
Cnemidophorus 312, 375
exsanguis 465
gularis 381
lemniscatus 380
sexlineatus 381–382
tigris 101, 267, 382
Coccoloba uvifera 316
Cocos 246
Coix lacryma-jobi 250
Coleonyx variegatus 465, 487
Colostethus 177
inguinalis 178
trinitatis 178
Coluber 191, 197, 200, 203, 205
constrictor 24–25, 259, 294
sp. 294
triaspis 202
Coniophanes 194, 197
bipunctatus 200
fissidens 195, 200
imperialis 195, 200
lateralis 200
picivittatus 200
quinquevittatus 200
Constrictor constrictor 101
Coprosma repens 337
Corallus 239
caninus 387
Coronella 202
Crataegus sp. 57
Crinia sp. 236
Crisantophis 191–193, 198
Crocodylus acutus 255, 257
johnstoni 256
mindorensis 256
niloticus 254–257
novaeguineae 249, 251–257
porosus 249, 251, 253, 255–256
Crotalus 299
adamanteus 103–104
atrox 23, 26, 101–104, 292–293
basiliscus 24, 103
basiliscus oaxacus 104
catalinensis 103–104
cerastes 98, 101, 103
cerastes cercobombus 100, 104
cerastes laterorepens 104
durissus 98, 101, 103
durissus culminatus 104
durissus terrificus 102, 104
enyo 98, 101, 103
enyo enyo 104
enyo furvus 104
exsul 103
horridus 98–99, 101–104, 260
intermedius 103
intermedius gloydi 101, 104
lannomi 98–101, 103–104
lepidus 101, 103
lepidus klauberi 104
lepidus lepidus 104
mitchellii 98, 101, 103
mitchellii mitchellii 99, 104
mitchellii stephensi 104
molossus 103
molossus molossus 104
molossus nigrescens 104
polystictus 101–104
pricei 98, 101, 103
pricei pricei 100, 104
pusillus 103–104
ruber 24, 103
ruber ruber 104
scutulatus 101, 103
scutulatus scutulatus 104
stejnegeri 98, 100, 103–104
tigris 101, 103–104
tortugensis 101, 103–104
transversus 103–104
triseriatus 101, 103
triseriatus aquilus 102, 104
triseriatus armstrongi 101, 104
unicolor 98, 101, 103–104
vegrandis 98, 101, 103–104
viridis 103, 259–260, 262–263
viridis helleri 104
viridis viridis 104, 259–260, 294
willardi 98, 101, 103
willardi meridionalis 100, 104
willardi silus 101, 104
Crotaphopeltis 191
Cryophis 191–193, 195–198
Cryptobatrachus 167
Cryptoblepharus plagiocephalus 236
Cryptobranchus alleganiensis 106, 111, 113, 115
Cupriguanus 120
Cyperus 252
Deirochelys reticularia 52
Dendrobates 182, 224
auratus 177–179, 181
granuliferus 177–180
histrionicus 177–178, 180–181
pumilio 177–181
trivittatus 178, 181
Denisonia sp. 35
Dermochelys coriacea 79, 315
Deschampsia flexuosa 346

Desmognathus fuscus 105, 361–363, 365–367
ochrophaeus 361, 363, 367
quadramaculatus 357–360
Diadophis 196
punctatus 294
Dicamptodon ensatus 360
Digiostoma truncatum 5
Diplodactylus vittatus 488
Dipsosaurus dorsalis 101
Drymarchon 196, 205
corais couperi 22–25
Drymobius melanotropis 133
Dysoxylum spectabile 337
Echinochloa 251
stagnina 250
Echinostoma ilocanum 5
Egernia margaretae 236
Elaphe 191, 197–198
carinata 205
chlorosoma 202
flavirufa 204, 206
guttata 23–26, 205
moellendorffi 203–206
obsoleta 22–26, 203, 205, 262
obsoleta bairdi 23
oxycephala 205
phaescens 204, 206
quatuorlineata 203
radiata 205–206
rosaliae 204, 206
rufodorsata 205
scalaris 205
schrenckii 205
sp. 294
subocularis 204–206
taeniura 203
triaspis 200–202, 206
vulpina 22–26, 206
Eleutherodactylus 1, 4, 224
antillensis 132
auriculatus 275
chlorophenax 270, 272, 274–276, 278–279
condor 95
cornutus 94
devillei 94
ernesti 90, **91**–95
galdi 94
hypostenor 269–279
inoptatus 269–272, 274–279
kirklandi 95
nigrovittatus 94
nortoni 270, 272, 274–279
pantoni 277
parapelates **269**–270, 272–278
peruvianus 94
quaquaversus 94
ruthae 269–279
ruthae aporostegus 277
ruthae bothroboans 277
ruthae ruthae 277
ruthae tychathrous 277
unistrigatus 95
Emydoidea blandingi 40, 49–52, 55, 324
Emydura macquarii 301–306
Epicrates monensis 132
Eremias lugubris 267
Eretmochelys 130
Erica tetralix 406
Eridiphas 190–198
slevini 200
Eristicophis macmahoni 101
Espeletia 155
Eublepharis kuroiwae 482, 488
kuroiwae kuroiwae 482–487
kuroiwae orientalis 482
kuroiwae splendens 482
Eumeces copei 29
fasciatus 7–13, 458, 466
inexpectatus 7–13, 458, 466
laticeps 7–12, 458–460, 462–463, 465–466
sp. 11
Eunectes murinus 387
Eupsophus calcaratus 173–176
migueli 173–177
roseus 173–177
vittatus 173–176
Eurycea 439
aquatica 423–426, 428–433, 435, 437
bislineata 105–106, 111, 358, 423–424, 426, 428–437
bislineata bislineata 423–424, 433
bislineata cirrigera 423–425, 429, 433
bislineata major 423–425
bislineata rivicola 423–425, 429, 433
bislineata wilderae 423–425, 432–434
cirrigera 423, 426, 429–431, 433–437
junaluska 423–426, 428–433, 435, 437
longicauda 424, 426–431, 433, 435
nana 105
neotenes 105
quadridigitata 105
wilderae 423, 426–427, 429–431, 434–437
Eutrombicula 188
Fagus grandifolia 329
Farancia 196
Fasciola hepatica 1, 5–6
Festuca rubra 346
Ficus ampelas 483
virgata 483
Gastrotheca 173
cavia 141, 144–145, 163–164, 167, 169–172
espeletia 141, 144–151, **152**, 153–155, 160–163, 165–172
griswoldi 141–142, 144–145, 164–167, 169–172
litonedis 141, 144–150, **155**–157, 160, 165–167, 169–172
lojana 141, 158, 167
marsupiata 141–142, 144–145, 160, 164–167, 169–172
marsupiata ecuatoriensis 163
marsupiata lojana 141, 158–159, 167
monticola 141, 144–149, 151, 155, 157–159, 165–167, 169–172
orophylax 141, 144–148, 150, 159, 164–167, 169–171
ovifera 164
peruana 160

plumbea 141, 144–148, 150, 158–160, 165–167, 169–172
pseustes 141, 144–151, 153–159, **160**–163, 165–167, 169–172
psychrophila 141, 144–148, 151, 153, 155, 157, 162–167, 169–172
riobambae 141–142, 144–151, 153–154, 160, 162–167, 169–172
ruizi 141, 144–152, 154, 162, 164–167, 169–172
testudinea 143–145, 164–165, 167, 169–172
weinlandii 143–145, 164–165, 167, 169–172
Gehyla variegata 488
Gerrhonotus coeruleus 29
Gonyosoma 197, 203, 205
Gopherus polyphemus 52
Graptemys geographica 79
pseudogeographica 79
pulchra 72, 79
Gyrinophilus porphyriticus 105, 358
Haideotriton 106, 113
wallacei 105
Halodule wrightii 317
Hanguana 251, 252
melayana 250
Helicops martii callostictus 387
pastazae 387
Helioporus sp. 226
Heloderma suspectum 101
Hemidactylium scutatum 105–107, 109, 111, 113, 116
Hemidactylus frenatus 487
Hemiphractus 167
Himantodes 190
Hippomane mancinella 316
Hydraticus 227, 230–231
bimarginata 229
Hydrochara sp. 229
Hydrops martii callostictus 387
Hyla argenteovirens 141
cinerea 224, 226–267
crucifer 216, 370–371
garbei 388
marsupiata 141
microcephala 479
pulchrilineata 243
quitoe 163
riobambae 141, 163
rubra 388
squirella 478
versicolor 479
Hylorana 96
Hynobius arisanensis 117–118
boulengeri 117–118
formosanus 117
kimurae 118
lichenatus 118
naevius 118
nebulosus 118
retardatus 118
sonani 117
stejnegeri 106–107, 111, 118
tsuensis 106–107, 111, 113
Hyperolius marmoratus 478
Hypnale 101
Hypsiglena 190, 193–197
affinis 192
latifasciata 191–192, 198
ochrorhynchus 192
torquata 192, 200
Hypsirhynchus 246
Ialtris 241
Iguana iguana 132
Imantodes 192–194, 197–198
cenchoa 195, 200, 387
gemmistratus 195
inornatus 133
lentiferus 195
Ipomea pes-caprae 319
Lacerta agilis 465
vivipara 345–347, 351, 405–408, 411–413
Lampropeltis 197–198, 201–202, 204–206
alterna 23
calligaster rhombomaculata 24–25
getulus 15, 24, 203
getulus californiae 25
getulus getulus 25
getulus holbrooki 15, 17–26
mexicana 22–24
mexicana alterna 23, 25
mexicana campbelli 23, 25
mexicana mexicana 23, 25
mexicana thayeri 23, 25
sp. 294
triangulum 22, 24
triangulum annulata 24
triangulum hondurensis 23, 25
triangulum sinaloae 23–26
triangulum syspila 24
Lampropholis guichenoti 236
Leersia 250
Leiocephalus 241, 246
eremitus 132
lunatus 243
melanochlorus 243–245
personatus 243
schreibersi 243
sp. 243
vinculum 243–244
Leiolopisma entrecasteauxi 35
sp. 236
trilineatum 236
weeksae 35
Leiopelma 128
Lepidoblepharis festae 386
Lepidophyma 281
flavimaculatum 280, 282–291
gaigae 282–289, 291
micropholis 282–288, 291
occulor 282–288, 291
pajapanensis 282–289, 291
smithii 282–288, 291
sylvaticum 282–288, 291
tuxtlae 282–288, 291
Leptodactylus 224
Leptodeira 192
annulata 195–196, 200
discolor 191
frenata 195–197, 200

guilleni 196
latifasciata 190–191, 195–198
nigrofasciata 193, 196–197, 200
punctata 195, 200
septentrionalis 194–196, 200
splendida 196
Leptodira 190
Leptophis 201
Libellula 231
vibrans 229
Limnodynastes dumerilii 236
ornatus 236
sp. 236
spenceri 236
tasmaniensis 236
Liophis breviceps 387
cobella 387
Liriodendron tulipifera 329
Litoria aurea 236
ewingii 236
moorei 236–237
sp. 236
Lygosoma 35
Lymnaea rubiginosa 4–6
Maaesa tenera 483
Mabuya bistriata 386
heathi 35
mabouya mabouya 386
Macaranga tanarius 483
Macropiper excelsum 337
Malaclemys terrapin 55
Masticophis 197
Meizodon 202
Melanoides tuberculata 5
Melia azedarach 483
Melicytus ramiflorus 337
Mephitis mephitis 61
Mesotriton 447, 455
Microtermis 5
Micrurus albicinctus 388
annellatus 388
browni 197
distans 197
filiformis subtilis 388
langsdorffi ornatissimus 388
laticollaris 197
lemniscatus helleri 388
Molinia caerulea 406
Moloch 267–268
Mus musculus 236–237
Myoporum laetum 337
Nardus stricta 346
Narthecium ossifragum 406
Nasua nasua 255
Natrix fasciata 294
natrix 294
sipedon 294
Necturus beyeri 106, 113, 115, 495
lewisi 490
punctatus 490–495
Neobatrachus pictus 236
sudelli 236
Nerodia 196
rhombifera 312
sipedon sipedon 294
Ninia hudsoni 388
Notechis 234, 237, 239
ater 233
scutatus 233, 236
Notonecta 230–231
sp. 229
Notophthalmus viridescens 370–371
Nototrema bolivianum 141
peruanum 141
plumbeum 141, 159
Nuphar variegatum 468
Ololygon 388
Opheodrys aestivus 23–25
Ophiophagus 24
Ophisaurus attenuatus 415–417
attenuatus attenuatus 415, 419, 422
attenuatus longicaudus 415–422
compressus 415–422
mimicus 415–**416**, 417–422
ventralis 415, 419–420, 422
Osteopilus dominicensis 243–244, 246–247
spp. 246
Oxybelis 246
brevirostris 133
Oxyrhopus formosus 387
occipitalis 387
Pachypalaminus 117–118
Pachyptila turtur 336
Pachytriton 115
Palaeotriton 447, 455
Parathelphusa convexia 4–6
Paratrophis banksii 337
Pelecanoides urinatrix 336
Peltocephalus dumeriliana 387
Philautus leucorhinus 479
Philodryas viridissimus 387
Pholidobolus 188
Phragmites 251–252
australis 302
karka 250
Phrynops gibbus 387
Phrynosoma 267–268
Phyllobates 182
lugubris 177–178, 180–181
Phyllodactylus marmoratus 236
Physalaemus 224
pustulosus 214, 478
Pinus palustris 421
sylvestris 406
Pituophis 24, 197, 201–202, 204–206
melanoleucus 25, 203, 262
melanoleucus catenifer 294
Plethodon 454
cinereus 105, 361, 365–367, 452
dorsalis 105
glutinosus 105
serratus 433
Pleurodeles 115
Podocnemis expansa 387
sextuberculata 387
unifilis 387
Polychrus marmoratus 386
Polygonum orientale 250
Prionodactylus 188
Pristidactylus achalensis 120–126

Procyon lotor 46, 55, 61, 255–256
Pseudacris crucifer 216–222
triseriata 231
Pseudemys scripta 50, 302, 305, 307, 309–313, 334
Pseudobranchus striatus 106, 111, 113, 115
Pseudoleptodeira 190–192, 194, 196
latifasciata 193, 195, 197–198, 200
Pseudomys albocinereus 236
Pseudophryne bibronii 236
guentheri 236
Pseudoraphis spinescens 250
Pseudotriton montanus 105
ruber 105
Pseustes poecilonotus polylepis 387
Psychotria manillensis 483
Pteridium aqualinum 421
Ptychoglossus 188
Puffinus gavia 337
griseus 337
Python 23
molurus 26, 294
Quercus alba 329
laevis 421
rubra 329
Rana boulengeri 97
cancrivora 1–6
catesbeiana 467, 478
clamitans 467–468, 477–478
pipiens 212
ridibunda 478
spinosa 97
sylvatica 370–371
temporaria 221
tiannanensis 97
virgatipes 467–469, 477–479
Ranidella signifera 236
Rattus fuscipes 236
sp. 236
Regina septemvittata 25
Rhabdophis 196
Rhadinaea 387
Rhadinophanes 191–193, 198
Rhagoletis 433
Rhinobothryum lentiginosum 387
Rhinocheilus 202
Rhinophis philippinus 101
Rhododendron spp. 357
Rhopalostylis sapida 337
Rhyacotriton olympicus 106, 111
Saccharum 251, 252
robustum 250
Salamandra bislineata 423
cirrigera 423–424
Salix spp. 302
Scaptochelys agassizi 334
Sceloporus 312
bicanthalis 35
clarki 126
cyanogenys 29
formosus 29
graciosus 465
grammicus 29, 36
jarrovi 29, 34
mucronatus 29
occidentalis 101
poinsetti 29, 34
undulatus 34, 36, 126
Scirpus suberminalis 468
Senticolis 200, **202**, 204
triaspis 201–203, 205–206
triaspis intermedia 202
triaspis mutabilis 202
triaspis triaspis 202
Seps 35
Sibon fischeri 133
Siphlophis cervinus 387
Siren intermedia 106, 111, 113, 115
sp. 492
Sistrurus catenatus 101, 103
catenatus edwardsi 101, 104
catenatus tergeminus 104
miliarius 98, 101, 103–104
miliarius miliarius 99
miliarius streckeri 99–100, 104
ravus 101, 103
ravus exiguus 104
ravus ravus 102, 104
Spalerosophis 203
Sphaerodactylus 247
becki 132
elegans 243
semasiops 133
Sphenodon punctatus 336
Sphenomorphus quoyii 236
Sporobolus virginicus 317, 319
Stefania 167
Stellastchasmus formosanum 5
Stereochilus marginatus 105
Sternotherus carinatus 71
minor 66–72, 312
minor minor 66
odoratus 71–72, 82–83, 86–89
Stilosoma 202
Storeria dekayi 24–25
Sus scrofa 254
Suta sp. 35
Syringodium filiforme 317
Tantalophis 191–193, 198
Tarenna gracilipes 483
Taricha 372
Telmatobius halli 208–211, 213
hauthali laticeps 214
marmoratus 208–213
montanus 208–210, 212, 215
pefauri 208–211, 213
peruvianus 208–211, 213
zapahuirensis 208–211
Terrapene carolina 328–334
coahuila 334
ornata 328–329, 333–334
Thalassia testudinum 317
Thamnodynastes pallidus 387
Thamnophis 24, 26, 197
brachystoma 294
elegans 259, 294
elegans atratus 294
melanogaster 25
proximus 294
radix haydeni 294
sauritus 229, 294

sirtalis 23, 25, 35, 294
sirtalis parietalis 259–260, 262
sp. 294
Thecadactylus rapicauda 132
Thoracostachyum 251–252
sumatranum 250
Trimorphodon 190–191, 194–195, 197
Trionyx muticus 50
Triturus 448
alpestris 446–447, 449–454, 457
boscai 446–447, 449–454, 457
cristatus 446–447, 449–454, 457
helveticus 446–447, 449–454, 457
italicus 446–447, 449–454, 457
marmoratus 446–447, 449–454, 457
montandoni 446–447, 449–454, 457
vittatus 446–447, 449–454, 457
vittatus ophryticus 447
vulgaris 106, 113, 115, 446–447, 449–454, 457
Tropidophis 132
Turpinia ternata 483
Typhlomolge rathbuni 105
Typhlops brongersmianus 387
minuisquamus 387
reticulatus 387
Typhlosaurus 267
Typhlotriton spelaeus 105–106, 111, 115
Uma notata 101
Uperoleia rugosa 478
Uracentron azureum 387
guentheri 387
Uromacer catesbyi 241–247
frenatus 241–247
frenatus dorsalis 246
oxyrhynchus 241–247
Ursus americanus 255
Utricularia spp. 468
Varanus spp. 255
Vipera berus 24, 101
lebetina schweizeri 101
Vulpes fulva 46
Xantusia bolsonae 288
henshawi 280, 284, 286–288, 487
henshawi bolsonae 280, 282–283, 286–288, 292
henshawi gracilis 287
henshawi henshawi 282–291
riversiana 282–288, 292
vigilis 29, 35, 280, 284, 286–289
vigilis arizonae 282–288, 292
vigilis extorris 280, 282–288, 292
vigilis sierrae 282–288, 292
vigilis wigginsi 282–283, 285–288, 292
vigilis vigilis 282–289, 292
Xenoboa sp. 101
Xenopus laevis 126

TOPICAL INDEX

HERPETOLOGICA

Volume 43 December 1987

ACTIVITY, *Eublepharis kuroiwae kuroiwae* (seasonal, daily, and diel) 482–489.

AGE STRUCTURE, *Pseudacris crucifer* (through skeletochronology) 216–223.

ALLOZYMES, *Eurycea bislineata* (geographic genetic variation) 423–446; *Gastrotheca* (14 nominal spp. compared) 141–173; *Triturus* (9 spp.) 446–457; Xantusiidae (3 spp. of *Xantusia* and 8 spp. of *Lepidophyma*) 280–292.

ANATOMY, *Eumeces laticeps* (cloaca and urodaeal gland) 458–466; *Hemidactylium scutatum* (female cloaca) 105–116.

BEHAVIOR, *Ameiva exsul* (activity budgets) 374–383; *Chrysemys picta marginata* (nesting activity in SW Quebec) 55–65; *Desmognathus fuscus* (brooding tenacity related to parental investment) 361–368; *Emydura macquarii* (atmospheric and aquatic basking) 301–306; *Eublepharis kuroiwae kuroiwae* (seasonal, daily, and diel activity) 482–489; *Eumeces fasciatus*, *E. inexpectatus*, and *E. laticeps* (agonistic between conspecific males) 7–14; *Lacerta vivipara* (thermoregulatory) 405–415; *Lampropeltis getulus holbrooki* (motor patterns of courtship and mating) 15–28; *Rana virgatipes* (vocal behavior and acoustic interactions) 467–481; *Telmatobius* (vocal behavior of 6 Chilean spp.) 208–216.

BIOLOGICAL CONTROL, *Rana cancrivora* (of pest crab and snail in ricefields and fish ponds of West Java) 1–6.

CLOACA, *Eumeces laticeps* (anatomy and urodaeal gland) 458–466; *Hemidactylium scutatum* (compared with other salamanders) 105–116.

COMPUTER, computer–microscope interface for measurements 384–385.

DEMOGRAPHY, *Lacerta vivipara* (3 populations compared in France) 345–356; *Necturus punctatus* 490–496.

DIET, *Necturus punctatus* 490–496; *Notechis* (differences between mainland and islands) 233–240; *Rana cancrivora* (in West Java) 1–6; *Uromacer* (3 spp. on Hispaniola) 241–248.

DISTRIBUTION MAPS, *Anadia bitaeniata* species group (5 spp. in Colombia and Venezuela) 184; *Barisia imbricata* (4 ssp. in Mexico) 30; 3 spp. of *Eleutherodactylus* (in southern Hispaniola) 275; 3 spp. of *Eurycea bislineata* complex 435; *Gastrotheca* (7 spp. in Ecuador, Colombia, and Peru) 154–155; *Ophisaurus mimicus* (SE United States) 429; *Senticolis triaspis* 201.

DOMINANCE, in *Ameiva exsul* (related to size) 374–383.

ECONOMIC IMPORTANCE, *Rana cancrivora* (as human food and ricefield pest control) 1–6.

EGGS, *Chelydra serpentina* (clutch and egg size) 39–54; *Chrysemys picta* (effects of hydric and thermal experimental environments on incubation) 393–404; *Sternotherus minor minor* (number and characteristics) 66–73.

ELECTROPHORESIS, *Eurycea bislineata* (geographic genetic variation) 423–446; *Gastrotheca* (relationships among 14 nominal spp.) 141–173; *Triturus* (relationships among 9 spp.) 446–457.

EMBRYOS, *Ambystoma maculatum* and *A. opacum* (mortality, field estimates and causes) 496–501.

GENETICS, *Crotalus atrox* (anomalous scutellation and pigmentation) 292–300.

HABITAT, *Necturus punctatus* 490–496; *Sphenodon punctatus* (burrow use, sharing with birds) 336–344; *Terrapene carolina* (in Indiana) 328–335.

HATCHLINGS, *Caretta caretta* (energetics of nest emergence and hatchling frenzy) 307–315; *Chelydra serpentina* (emergence dates and survival) 39–54; *Chrysemys picta* (effects of hydric and thermal experimental environments during incubation) 393–404; *Chrysemys picta marginata* (emergence, predation, survival in SW Quebec) 55–65; *Sternotherus minor minor* (size in C Florida) 66–73.

HOME RANGE, *Ameiva exsul* (influenced by sex and body size) 374–383.

HOMING, *Terrapene carolina* (in Indiana) 328–335.

KARYOTYPES, *Dendrobates* (4 spp.) and *Phyllobates* (1 sp.) 177–182; *Hynobius arisanensis* (compared with Japanese spp.) 117–119; *Pristidactylus achalensis* (polymorphisms of microchromosomes and nucleolar organizer region) 120–127.

LOCOMOTION, *Caretta caretta* (locomotor performance of hatchlings) 307–315.

LONGEVITY, *Terrapene carolina* (in Indiana) 328–335.

MATING, *Lampropeltis getulus holbooki* (analysis of behavioral motor patterns) 15–28.

MORPHOLOGY, *Bombina orientalis* (computer-assisted measurements) 384–385; *Crotalus atrox* (inheritance of anomalous scutellation and pigmentation in a lab colony) 292–300; *Crotalus* and *Sistrurus* (dorsal scale microdermatoglyphic patterns of 30 spp.) 98–104; *Notechis* (mainland and island differences in body size) 233–240; *Pseudoleptodeira latifasciata* (compared with other cat-eyed snakes and allies) 190–200; *Uromacer* (body morphometrics related to diet in 3 spp.) 241–248.

MOVEMENTS, *Terrapene carolina* (in Indiana) 328–335.

NESTS, *Caretta caretta* (emergence from in Florida) 307–315; *Chelonia mydas* (temperatures at Tortuguero, Costa Rica) 74–81; *Chelydra serpentina* (SE Michigan) 39–54; *Chrysemys picta marginata* (SW Quebec) 55–65; *Crocodylus novaeguineae* and *C. porosus* in Papua New Guinea (descriptions, locations, season) 249–258; *Dermochelys coriacea* (placement and fate, U.S. Virgin Islands) 315–323.

NUMERICAL TAXONOMY, *Eupsophus* (identification of 4 Chilean spp. through principal components analysis of body dimensions) 173–177.
PALATABILITY, *Bufo americanus* tadpoles (to dragonfly naiads, giant water bugs, and newts) 369–373.
PHEROMONES, *Eumeces fasciatus* group (involved in conspecific male agonistic behavior) 7–14; *Eumeces laticeps* (pheromone-secreting urodaeal gland in cloaca) 458–466.
PHYLOGENY, *Crotalus* and *Sistrurus* (cladistic analysis of dorsal scale microdermatoglyphic patterns of 30 spp.) 98–104; *Gastrotheca* (of 14 nominal spp., based on electromorphic data) 141–173; Plethodontidae (based on female cloacal anatomy) 105–116; *Pseudoleptodeira latifasciata* and allied genera (based on albumin immunological distances) 190–200; *Triturus* (among 9 spp., based on electromorphic data) 446–457; Xantusiidae (relationships among 20 samples of 11 spp., based on electrophoretic analysis) 280–292.
PHYSIOLOGY, *Caretta caretta* (anaerobic metabolism during nest emergence and hatchling frenzy) 307–315.
POPULATION DENSITY, *Desmognathus quadramaculatus* (experimentally related to rock density in North Carolina) 357–361; *Lacerta vivipara* (3 populations compared in France) 345–356; *Sphenodon punctatus* 336–344; *Terrapene carolina* (in Indiana) 328–335.
POPULATION DYNAMICS, *Lacerta vivipara* (3 populations in France) 345–356.
PREDATION, on *Bufo americanus* tadpoles (antipredator mechanisms) 369–373; *Chrysemys picta marginata* (nest mortality by raccoons in SW Quebec) 55–65; by predatory insects on tadpoles (related to pool size) 224–232.
REPRODUCTION, *Barisia imbricata* (gonadal cycles, seasonal activity, gestation, clutch size, neonates) 29–38; *Chelydra serpentina* (nesting ecology, clutch size and frequency, hatching, demography and survivorship, nest depredation in SE Michigan) 39–54; *Chelydra serpentina* (predicting onset of nesting season in NC Ontario) 324–328; *Chrysemys picta marginata* (nesting season and hatching success in SW Quebec) 55–65; *Crocodylus novaeguineae* and *C. porosus* (breeding size and season, clutches, reproductive success in Papua New Guinea) 249–258; *Lacerta vivipara* (3 populations compared in France) 345–356; *Sternotherus minor minor* (maturation, vitellogenesis, oviposition, and clutch size and frequency in C Florida) 66–73; *Sternotherus odoratus* (experimental effects of temperature and photoperiod on the ovarian cycle) 82–90.
SEROLOGY, albumin immunological distances among some cateyed snakes 190–200.
SEX DETERMINATION, *Chelonia mydas* (temperature dependent on Tortuguero beach, Costa Rica) 74–81.
SEX RATIO, *Chelonia mydas* (effects of nest temperature at Tortuguero, Costa Rica) 74–81; *Notechis* (mainland and island populations) 233–240.
SONOGRAMS, 5 spp. of *Eleutherodactylus inoptatus* group (Dominican Republic and Haiti) 277; *Rana virgatipes* (vocal repertoire) 469; *Telmatobius* (contact and distress calls) 208–216.
SPERMATHECA, *Hemidactylium scutatum* (compared with other salamanders) 105–116.
SURVIVORSHIP, *Ambystoma maculatum* (embryos and juveniles in North Carolina) 496–501; *Chelydra serpentina* (hatchlings in Michigan) 39–54; *Terrapene carolina* (in Indiana) 328–335.
SYSTEMATICS, *Amolops macrorhynchus* (new sp. from China) 95–97; *Anadia bitaeniata* species group (5 spp. from Colombia and Venezuela, 1 new) 182–190; *Eleutherodactylus ernesti* (new sp. from Ecuador) 90–95; *Eleutherodactylus parapelatus* (new sp. from Haiti) and *E. inoptatus* group 269–279; *Eurycea bislineata* (comprises at least 3 spp.) 423–446; *Gastrotheca* (data on allozymes, morphometrics, structure, and coloration of 9 spp., 3 new) 141–173; *Hynobius arisanensis* (karyotype compared with Japanese spp.) 117–119; *Ophisaurus mimicus* (new sp. from SE United States) 414–423; *Pseudoleptodeira* (status and relationships) 190–200; *Senticolis* (new genus for *Elaphe triaspis*, compared with other Lampropeltiini) 200–207; *Triturus* (allozyme data on 9 spp.) 446–457; Xantusiidae (allozyme evolution and phylogeny among 11 spp.) 280–292.
THERMOREGULATION, *Crotalus v. viridis* (temperature drop in solitary and aggregated snakes compared) 259–264; *Emydura macquarii* (through atmospheric and aquatic basking) 301–306; *Lacerta vivipara* (seasonal) 405–415.
UNISEXUALITY, *Lepidophyma flavimaculatum* (1 population) 280–292.
VIVIPARITY, *Barisia imbricata* (in high elevation Mexico) 29–38.
VOCALIZATIONS, *Pseudacris crucifer* (age, size, and call frequency) 216–223; *Rana virgatipes* (and acoustic interactions) 467–481; *Telmatobius* (6 Chilean spp.) 208–216.